

Number of births



Application for Data for Research Purposes from Perinatal Services BC Appendix A: BCPDR Data Fields Checklist

Pro	ject Title	Click here to enter text.	Applies	Click here to enter ext.	
Date	Range	Click here to enter text.			
Othe	r date range criteria	Click here to enter text.			
ВС	Perinatal Data Re	egistry (BCPDR)			
	ril 1, 2000 to Marc	• , ,			
captur Trans For de BCPD For al ensur • L					
	Additional Exclusions : Therapeutic abortions and records with "Discharge Disposition" to Adoption or Foster Care are excluded from the BCPDR facilitated by the WHRI.				
	Mother Delivery Episode of Care Information (April 1, 2000 to March 31, 2018)				
NOTE will ha	Delivery records include deliveries in acute care facilities and deliveries at home attended by registered midwives. NOTE: Until March 31, 2014, a woman who delivered at home with a registered midwife who was admitted to acute care within 24 hours of delivery will have the acute care admission as her Delivery record (total record count=1). Effective April 1, 2014, a woman who delivered at home with a registered midwife and who was admitted to acute care within 24 hours of delivery will have a home Delivery record and a subsequent Postpartum acute care record (total record count=2).				
			The Health Authority (HA) of mother's usual residence as determined by resident postal code.		
	Baby sequence		Sequence of baby in the current pregnancy.		
	Number of hirths		Total number of babies delivered in the current		

pregnancy.

Version: 2 Last Modified: 2019/11/25

Mother's date of birth – Year		
Mother's date of birth – Month		
Mother's date of birth – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers.		
Mother age at delivery – **Research rationale describing why this field is required must be supplied in your application before it will be considered for release:	Mother's age (in years) calculated at date of delivery.	
Click here to enter text.	iviolitiei s age (iii years) calculated at date of delivery.	
Place of delivery – Institution number – Replaced by project-specific identification number	Location where mother received care.	
Institution from – Replaced by project-specific identification number	Institution from which mother arrived to the current episode of care.	
Mother transferred in	A flag (i.e., Yes, Null) to indicate the mother was transferred in from another acute care institution for the delivery episode of care.	
Institution to – Replaced by project-specific identification number	Institution to which mother was transferred from the current episode of care.	
Mother transferred out	A flag (i.e., Yes, Null) to indicate the mother was transferred out to another institution after delivery.	
Mother transferred up	A flag (i.e., Yes, Null) to indicate the mother with delivery episode was transferred directly from delivery hospital to a hospital with a higher level of care.	
Admission date – Year	Year mother was admitted for the current episode of care.	
Admission date – Month	Month mother was admitted for the current episode of care.	
Admission date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers.	Day mother was admitted for the current episode of care.	
Discharge date – Year	Year mother was discharged from the current episode of care.	
Discharge date – Month	Month mother was discharged from the current episode of care.	
Discharge date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers.	Day mother was discharged from the current episode of care.	
Fiscal year	The fiscal year during which the mother was discharged (i.e., Apr 1 to Mar 31).	
Total length of stay	Total length of stay (in hours) for the delivery hospitalization. Note: Always null for deliveries at home.	
Antepartum length of stay	Time, in hours, between admission to delivery episode and delivery of the first baby. Note: Always	

		null for deliveries at home.	
	Postpartum length of stay	Time, in hours, between delivery of the placenta and discharge from the episode of care. Note: Always null for deliveries at home.	
Past	Obstetric History		
	Gravida	Total number of prior plus current pregnancies.	
	Parity	Indicates whether woman has previously delivered a pregnancy that reached 20 weeks gestation or 500 grams birth weight (i.e., multiparous, nulliparous, unknown).	
	Number of previous term deliveries	Total number of previous pregnancies delivered at ≥ 37 completed weeks gestation.	
	Number of previous preterm deliveries	Total number of previous pregnancies delivered between 20 to 36 completed weeks gestation.	
	Number of living children	Total number of children the mother has given birth to, who are currently living.	
	Number of previous spontaneous abortions	Total number of previous natural or spontaneous losses in pregnancy < 20 completed weeks and < 500 grams.	
	Number of previous cesarean sections	Total number of previous pregnancies resulting in a cesarean delivery ≥ 20 completed weeks gestation.	
	Number of previous vaginal deliveries	Total number of previous pregnancies resulting in a vaginal delivery ≥ 20 completed weeks gestation.	
	Prior neonatal death	A flag (i.e., Yes, Null) to indicate mother had at least one prior live born infant, who died within the first 28 days of life.	
	Prior stillbirth	A flag (i.e., Yes, Null) to indicate mother had at least one prior stillbirth or intrauterine death documented.	
	Prior low birthweight baby	A flag (i.e., Yes, Null) to indicate mother had at least one prior low birth weight baby (< 2,500 g) at ≥ 20 weeks gestation.	
	Prior macrosomic baby	A flag (i.e., Yes, Null) to indicate mother had at least one prior macrosomic baby (birth weight > 4,000g).	
	Rh isoimmunization (past pregnancy)	A flag (i.e., Yes, Null) to indicate mother had a previous pregnancy in which isoimmunization occurred.	
	Major congenital anomalies (past pregnancy)	A flag (i.e., Yes, Null) to indicate mother had at least one previous pregnancy in which the baby or fetus displayed a major congenital anomaly.	
	History of mental illness – Any	A flag (i.e., Yes, Null) to indicate any history of mental illness (depression, previous postpartum depression, anxiety, bipolar disorder, other, or unknown type) prior to or during the current pregnancy.	2008/09 onwards
	History of mental illness – Anxiety	A flag (i.e., Yes, Null) to indicate mother has documented history of anxiety.	2008/09 onwards
	History of mental illness – Depression	A flag (i.e., Yes, Null) to indicate mother has	2008/09 onwards

		documented history of depression.	
	History of mental illness – Bipolar	A flag (i.e., Yes, Null) to indicate mother has documented history of bipolar disorder.	2008/09 onwards
	History of mental illness – Postpartum depression (past pregnancy)	A flag (i.e., Yes, Null) to indicate mother has documented history of postpartum depression.	2008/09 onwards
	History of mental illness – Other	A flag (i.e., Yes, Null) to indicate mother has documented history of other mental illness.	2008/09 onwards
	History of mental illness – Unknown	A flag (i.e., Yes, Null) to indicate mother has documented history of mental illness, type unspecified.	2008/09 onwards
Curr	ent Pregnancy		
	First contact with physician/midwife date – Year	Year of mother's first contact with a physician/midwife for this pregnancy.	
	First contact with physician/midwife date – Month	Month of mother's first contact with a physician/midwife for this pregnancy.	
	First contact with physician/midwife date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day of mother's first contact with a physician/midwife for this pregnancy.	
	Number of antenatal visits	Total number of antenatal visits with the primary care provider.	
	Total antenatal hospital admissions (prior to delivery admission)	Total prior inpatient hospital admissions, to any facility, for any reason, during the current pregnancy (excluding current delivery admission).	
	Pre-pregnancy weight	Mother's weight (in kilograms) before pregnancy or ≤ 11 weeks completed gestation. Note: Approx. 21% missing.	
	Admission weight	Mother's weight (kg) at the time of admission for delivery, or the last weight documented ≤ 7 days prior to delivery. Note: Approx. 29% missing.	
	Weight gain in pregnancy	Mother's weight gain (in kilograms) during the pregnancy. Note: Approx. 39% missing.	
	Height	Mother's height (in cm). Note: Approx. 20% missing.	
	Body Mass Index (BMI)	Body Mass Index number of the mother, based on pre-pregnancy weight. Note: Approx. 30% missing.	
	Body Mass Index (BMI) group	Body mass index category of the mother, based on pre-pregnancy weight (e.g., underweight, normal, overweight).	
	Lone Parent	A flag (i.e., Yes, No, Unknown) to indicate lone parent status.	2000/01-2007/08
	Blood type	Classification of mother's blood type (e.g., A+, AB-).	
	Last Normal Menstrual Period (LNMP) date – Year	Year of LNMP. Note: Approx. 24% missing.	
	Last Normal Menstrual Period (LNMP) date – Month	Month of LNMP. Note: Approx. 24% missing.	
	Last Normal Menstrual Period (LNMP) date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day of LNMP. Note: Approx. 24% missing.	

First ultrasound date – Year	Year of first ultrasound (< 20 weeks). Note: Approx. 28% missing.	
First ultrasound date – Month	Month of first ultrasound (< 20 weeks). Note: Approx. 28% missing.	
First ultrasound date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day of first ultrasound (< 20 weeks). Note: Approx. 28% missing.	
Gestational age at first ultrasound – completed weeks	Gestational age, in weeks, when the first ultrasound (<20 weeks) was performed.	
Gestational age at first ultrasound – days	Gestational age, in days, when the first ultrasound (<20 weeks) was performed.	2008/09 onwards
Gestational age at delivery (by LNMP)	Gestational age at delivery in completed weeks (calculated by Last Normal Menstrual Period; LNMP). Note: Checked automatically because this field should always be requested when delivery or newborn records requested.	
Gestational age at delivery (by first ultrasound date)	Gestational age at delivery in completed weeks (calculated by first ultrasound date). Note: Checked automatically because this field should always be requested when delivery or newborn records requested.	
Gestational age at delivery (by algorithm)	Gestational age at delivery in completed weeks, calculated by algorithm incorporating LNMP, first ultrasound, newborn examination, and maternal chart. Note 1: Algorithms updated in 2013. Note 2: Checked automatically because this field should always be requested when delivery or newborn records requested.	
Rh immunoglobulin given, earliest date antepartum – Year	Year of the first antepartum injection of Rh immunoglobulin administered to an Rh-negative mother during the current pregnancy.	
Rh immunoglobulin given, earliest date antepartum – Month	Month of the first antepartum injection of Rh immunoglobulin administered to an Rh-negative mother during the current pregnancy.	
Rh immunoglobulin given, earliest date antepartum – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day of the first antepartum injection of Rh immunoglobulin administered to an Rh-negative mother during the current pregnancy.	
Hemoglobin level third trimester	Lowest hemoglobin value for the third trimester.	
Bleeding (<20 weeks)	A flag (i.e., Yes, Null) to indicate mother had antepartum bleeding in pregnancy < 20 weeks gestation.	
Antepartum hemorrhage (≥20 weeks)	A flag (i.e., Yes, Null) to indicate mother had antepartum hemorrhage or bleeding in pregnancy ≥ 20 weeks gestation, including bleeding from cervical polyps.	
Pregnancy induced hypertension	A flag (i.e., Yes, Null) to indicate care provider diagnosed mother with gestational hypertension during the current pregnancy.	
Proteinuria	A flag (i.e., Yes, Null) to indicate care provider diagnosed proteinuria.	

	1	ı
Rh blood antibodies	A flag (i.e., Yes, Null) to indicate mother developed or showed signs of Rh (anti-D) antibodies in her blood in the current pregnancy.	
Other blood antibodies	A flag (i.e., Yes, Null) to indicate mother developed or showed signs of antibodies in her blood, other than Rh antibodies, in the current pregnancy.	
Intrauterine growth restriction (IUGR) identified as risk during antenatal period	A flag (i.e., Yes, Null) to indicate health care provider identified intrauterine growth restriction (IUGR) during the antenatal period.	
Diabetes (any)	A flag (i.e., Yes, Null) to indicate pre-existing or gestational diabetes.	
Gestational diabetes (insulin dependent)	A flag (i.e., Yes, Null) to indicate gestational diabetes (insulin dependent).	
Gestational diabetes (non-insulin dependent)	A flag (i.e., Yes, Null) to indicate gestational diabetes (non-insulin dependent).	
Diabetes mellitus (insulin dependent)	A flag (i.e., Yes, Null) to indicate pre-existing diabetes mellitus Type 1 or Type 2, insulin used.	
Diabetes mellitus (non-insulin dependent)	A flag (i.e., Yes, Null) to indicate pre-existing diabetes mellitus Type 1 or Type 2, insulin not used.	
Abnormal glucose factor	A flag (i.e., Yes, Null) to indicate care provider diagnosed Abnormal Glucose Factor in pregnancy.	2000/01-2010/11
Hypertension (≥ 140/90)	A flag (i.e., Yes, Null) to indicate mother had a blood pressure reading of ≥ 140/90 on two consecutive readings during the pregnancy, prior to labour. NOTE: This variable alone is insufficient to identify women diagnosed with hypertension.	
Antihypertensive drugs	A flag (i.e., Yes, Null) to indicate mother received antihypertensive drugs during her pregnancy (antepartum period only).	
Hypertensive chronic renal disease	A flag (i.e., Yes, Null) to indicate mother had hypertension associated with chronic renal disease in the current pregnancy.	
Hypertension due to other causes	A flag (i.e., Yes, Null) to indicate mother had hypertension as a result of another cause during pregnancy, labour, or the postpartum period.	
Drug use during pregnancy identified as a risk	A flag (i.e., Yes, Null) to indicate care provider lists mother's use of drugs (prescription, non-prescription, illicit) as a risk factor in this pregnancy.	2000/01-2007/08
Substance use during pregnancy – Any	A flag (i.e., Yes, Null) to indicate mother used any of the following substances at any time during the current pregnancy: Heroin/opioids, cocaine, methadone, solvents, or marijuana; OR care provider lists use of prescription, 'other', or unknown other drug as a risk to the pregnancy.	2008/09 onwards
Substance use during pregnancy – Heroin/opioids	A flag (i.e., Yes, Null) to indicate heroin/opioid use during pregnancy, including before woman knew she was pregnant.	2008/09 onwards
Substance use during pregnancy – Cocaine	A flag (i.e., Yes, Null) to indicate cocaine use during pregnancy, including before woman knew she was	2008/09 onwards

	pregnant.	
Substance use during pregnancy – Methadone	A flag (i.e., Yes, Null) to indicate methadone use during pregnancy, including before woman knew she was pregnant.	2008/09 onwards
Substance use during pregnancy – Solvents	A flag (i.e., Yes, Null) to indicate solvent use during pregnancy, including before woman knew she was pregnant.	2008/09 onwards
Substance use during pregnancy – Prescription drugs	A flag (i.e., Yes, Null) to indicate use of a prescription drug is noted as a risk in the pregnancy.	2008/09 onwards
Substance use during pregnancy – Marijuana	A flag (i.e., Yes, Null) to indicate marijuana use during pregnancy, including before woman knew she was pregnant.	2008/09 onwards
Substance use during pregnancy – Other	A flag (i.e., Yes, Null) to indicate other substance use during pregnancy, including before woman knew she was pregnant.	2008/09 onwards
Substance use during pregnancy – Unknown	A flag (i.e., Yes, Null) to indicate care provider lists mother's use of an unspecified drug as a risk at any time during current pregnancy.	2008/09 onwards
No selected risks	A flag (i.e., Yes, Null) to indicate the mother did not have any of the specific risk factors collected in the PDR identified in the current pregnancy, past pregnancies, or in the mother's medical history.	
Alcohol during pregnancy identified as a risk	A flag (i.e., Yes, Null) to indicate care provider lists mother's use of alcohol as a risk factor in this pregnancy.	
T-ACE score	Final value of T-ACE questionnaire. Note: >99% null values.	2000/01-2007/08
TWEAK score	Final value of TWEAK questionnaire. Note: >96% null values	2008/09 onwards
Average number of alcoholic drinks per week	Average number of alcoholic drinks consumed per week by mother during current pregnancy.	2008/09 onwards
Binge drinking	Mother consumed ≥ 4 alcoholic drinks at one time during the current pregnancy.	2008/09 onwards
Smoking during current pregnancy	Mother smoked tobacco products during pregnancy.	
Cigarettes per day	Number of documented cigarettes smoked per day during pregnancy.	
Exposure to second hand smoke	Mother was regularly exposed to indoor smoke any time during the current pregnancy, either at home or work.	2008/09 onwards
HIV test done during pregnancy	HIV testing was performed during this pregnancy.	2004/05 onwards
Maternal serum screening offered during current pregnancy	Indicates whether maternal serum screening offered during current pregnancy (i.e., Yes, No, Unknown).	2004/05 onwards
Group B strep test done during current pregnancy	Mother had Group B Strep (GBS) testing done during current pregnancy (i.e., Yes, No, Unknown).	2004/05 onwards
Group B strep testing results	Indicates Group B strep testing results (i.e., positive, negative, unknown).	2004/05 onwards
HBsAg testing	Indicates Hepatitis B surface antigen (HBsAg) testing was performed at any time during the current pregnancy, prior to delivery (i.e., Yes, No, Unknown).	2008/09 onwards

	HBsAg testing results	Indicates HBsAg testing results (i.e., positive, negative, unknown).	2008/09 onwards
	In vitro fertilization used for current pregnancy	Indicates mother had in-vitro fertilization to achieve the current pregnancy. (i.e., Yes, No, Unknown).	2008/09 onwards
	School years completed	Total number of school years completed by the mother. Note: Approx. 80% Null values.	2004/05 onwards
Lab	our and delivery		
	Cervical dilation on admission	Measurement of cervical dilation, in centimeters, taken within the first hour of admission for the delivery episode.	
	Cervical dilation prior to cesarean	Last recorded measurement of cervical dilation during active labour, in centimeters, prior to cesarean delivery.	2008/09 onwards
	Rupture of membranes date – Year	Year of artificial/spontaneous rupture of the amniotic sac.	
	Rupture of membranes date – Month	Month of artificial/spontaneous rupture of the amniotic sac.	
	Rupture of membranes date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day of artificial/spontaneous rupture of the amniotic sac.	
	Length of time from rupture of membranes to first stage of labour (hours)	Hours between rupture of membranes and beginning of first stage of labour. Based on first infant delivered.	
	Start of first stage of labour date – Year	Year when there was onset of regular uterine contractions and cervical dilation.	
	Start of first stage of labour date – Month	Month when there was onset of regular uterine contractions and cervical dilation.	
	Start of first stage of labour date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day when there was onset of regular uterine contractions and cervical dilation.	
	Length of the first stage of labour	Duration of first stage of labour (rupture of membranes to full cervical dilation), in hours.	
	Start of second stage of labour date – Year	Year there was full cervical dilation and delivery of the newborn commenced.	
	Start of second stage of labour date – Month	Month there was full cervical dilation and delivery of the newborn commenced.	
	Start of second stage of labour date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day there was full cervical dilation and delivery of the newborn commenced.	
	Length of the second stage of labour	Duration of second stage of labour (full cervical dilation to delivery of infant), in hours.	
	Baby delivery date – Year	Year baby was delivered.	
	Baby delivery date – Month	Month baby was delivered.	
	Baby delivery date – Day– This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day baby was delivered.	

Placenta delivery date – Year	Year of placenta delivery date.	
Placenta delivery date – Month	Month of placenta delivery date.	
Placenta delivery date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day of placenta delivery date.	
Length of third stage of labour	Duration of third stage of labour (delivery of infant to delivery of placenta), in hours.	
Length of time from rupture of membranes to first baby delivery time	Time between rupture of membranes and delivery, in hours.	
Fetal surveillance during labour	Fetal surveillance during labour (i.e., external electronic fetal monitoring, internal electronic fetal monitoring, external and internal electronic monitoring, no labour, no electronic monitoring). Fetal surveillance during labour (i.e., auscultation only, auscultation and external electronic fetal monitoring, external electronic fetal monitoring only, internal electronic fetal monitoring only, auscultation and internal electronic fetal monitoring, external and internal electronic fetal monitoring, all, no labour, none).	2000/01-2003/04 2004/05-onwards
Labour initiation — spontaneous	A flag (i.e., Yes, Null) to indicate onset of regular contractions and progressive dilation of the cervix occurred without instrumental or medicinal assistance.	
Labour initiation — induced	A flag (i.e., Yes, Null) to indicate instrumental or medicinal assistance was used to initiate labour.	
Labour initiation — none	A flag (i.e., Yes, Null) to indicate woman did not labour.	
Labour initiation — unknown	A flag (i.e., Yes, Null) to indicate unknown how labour commenced.	
Labour type	Indicates labour type (i.e., spontaneous, induced, no labour, unknown)	
Labour induction – Artificial rupture of membranes (ARM)	A flag (i.e., Yes, Null) to indicate labour was induced using artificial rupture of membranes.	
Labour induction – Oxytocin	A flag (i.e., Yes, Null) to indicate labour was induced using oxytocin.	
Labour induction – Prostaglandin	A flag (i.e., Yes, Null) to indicate labour was induced using prostaglandin.	
Labour induction – Other agent	A flag (i.e., Yes, Null) to indicate labour was induced using another method.	
Primary indication for induction	Indicates the primary indication that an external agent was used to initiate labour (i.e., post-term, prelabour ROM, fetal compromise, other maternal condition, logistics, fetal demise, other, unknown, not applicable). Note that the following options were added in 2008/09: Hypertension in pregnancy, antepartum hemorrhage, chorioamnionitis, diabetes.	
Labour augmentation	A flag (i.e., Yes, Null) to indicate labour was augmented.	

Method of labour augmentation – Artificial rupture of membranes (ARM)	A flag (i.e., Yes, Null) to indicate labour was augmented using artificial rupture of membranes.	
Method of labour augmentation – Oxytocin	A flag (i.e., Yes, Null) to indicate labour was augmented using oxytocin.	
Method of labour augmentation – Other agent	A flag (i.e., Yes, Null) to indicate labour was augmented using another method.	
Method of labour augmentation – Prostaglandin	A flag (i.e., Yes, Null) to indicate labour was augmented using prostaglandin.	2000/01-2007/08
Anesthesia/analgesia during labour – None	A flag (i.e., Yes, Null) to indicate no anesthetic or analgesic was given during labour (first, second or third stage).	
Anesthesia/analgesia during labour – Entonox (nitronox)	A flag (i.e., Yes, Null) to indicate entonox (nitronox) anesthetic was given during labour (first, second or third stage).	
Anesthesia/analgesia during labour – Local	A flag (i.e., Yes, Null) to indicate local anesthetic was given during labour (first, second or third stage).	
Anesthesia/analgesia during labour – Pudendal	A flag (i.e., Yes, Null) to indicate pudendal anesthetic was given during labour (first, second or third stage).	
Anesthesia/analgesia during labour – Epidural	A flag (i.e., Yes, Null) to indicate epidural anesthetic was given during labour (first, second or third stage).	
Anesthesia/analgesia during labour – Spinal	A flag (i.e., Yes, Null) to indicate spinal anesthetic was given during labour (first, second or third stage).	
Anesthesia/analgesia during labour – General	A flag (i.e., Yes, Null) to indicate general anesthetic was given during labour (first, second or third stage).	
Anesthesia/analgesia during labour – Narcotics	A flag (i.e., Yes, Null) to indicate mother received narcotics during labour (first, second or third stage).	
Anesthesia/analgesia during labour – Other	A flag (i.e., Yes, Null) to indicate other anesthetic or analgesic was given during labour (first, second or third stage).	
Anesthesia/analgesia during labour – Unknown	A flag (i.e., Yes, Null) to indicate type of anesthetic or analgesic administered during labour (first, second or third stage) is unknown.	
Mode of delivery	Method of extraction/delivery of newborn from the mother (i.e., cesarean section, vaginal)	
Mode of delivery – detailed	Expanded classification of method of extraction/delivery of newborn from the mother (i.e., emergency primary, emergency repeat, elective primary, elective repeat, forceps and vacuum, forceps, vacuum, other instrument, spontaneous).	
Cesarean section type	Type of cesarean section (i.e., primary elective, primary emergent, repeat elective, repeat emergent).	
Cesarean section incision	Type of cesarean section incision.	
Primary indication for cesarean delivery	Primary/principal reason (indication) for cesarean delivery (i.e., breech, dystocia/CPD, non-reassuring fetal heart rate pattern, repeat cesarean, abruptio placenta, placenta previa, other, malposition/malpresentation, active herpes. Note that the following option was added in 2008/09: VBAC declined or maternal request)	
Vaginal birth after cesarean (VBAC) eligible	Mother is eligible to deliver this pregnancy by VBAC.	
Vaginal birth after cesarean (VBAC) attempted	Whether woman attempted a VBAC in this pregnancy.	
Vaginal birth after cesarean (VBAC) successful	Woman had a successful VBAC in this pregnancy.	
Deliverer provider type	The health care provider (or person) who physically delivers the baby. Note: not necessarily the same as the provider who was seen for antenatal care.	

	Baby position in labour	Position of baby's head relative to the birth canal during labour.
	Baby position at delivery	Position of baby's head relative to the birth canal at delivery.
	Baby presentation in labour	Part of the baby's body that is presenting in reference to the birth canal during labour.
	Baby presentation at delivery	Part of baby's body that is presenting in reference to the birth canal at the time of delivery.
	Obstetric trauma	A flag (i.e., Yes, Null) to indicate woman experienced obstetric trauma during the current delivery episode.
	Perineal trauma – Intact perineum	A flag (i.e., Yes, Null) to indicate perineum/vagina/cervix was intact.
	Perineal trauma – Unknown	A flag (i.e., Yes, Null) to indicate condition of the perineum/vagina/cervix is unknown.
	Perineal trauma – Episiotomy	A flag (i.e., Yes, Null) to indicate an episiotomy was done.
	Perineal trauma – Episiotomy type	Type of episiotomy performed (i.e., median, mediolateral).
	Perineal trauma – Laceration	Tear and/or rupture occurred to the vagina or perineum during delivery, excluding abrasions.
	Perineal trauma – Laceration degree	Highest degree of laceration sustained during delivery (i.e., 1-4).
	Perineal trauma – Cervical tear	A flag (i.e., Yes, Null) to indicate mother experienced cervical injury during delivery.
	Perineal trauma – Other tear	A flag (i.e., Yes, Null) to indicate another type of tear or laceration to the perineum was sustained during delivery.
Othe	er Episode of Care Information	
	Blood transfusion given	A flag (i.e., Yes, Null) to indicate mother received whole or packed red blood cells during this admission.
	Blood transfusion units – Number of units transfused antepartum	Total number of units of whole or packed red blood cells the mother received during the antepartum period of this pregnancy.
	Blood transfusion units – Number of units transfused intrapartum	Total number of units of whole or packed red blood cells the mother received during the intrapartum period of this pregnancy.
	Blood transfusion units – Number of units transfused postpartum	Total number of units of whole or packed red blood cells the mother received during the postpartum period of this pregnancy.
	Blood transfusion units – Total number of units transfused	Total number of units of whole or packed red blood cells the mother received during this pregnancy.
	Eligible for postpartum Rh immunoglobulin	Mother is eligible to receive Rh Immunoglobulin postpartum (i.e., Yes, No, Unknown).
	Date postpartum Rh immunoglobulin – Year	Year of the postpartum injection of Rh immunoglobulin during the delivery episode.
	Date postpartum Rh immunoglobulin – Month	Month of the postpartum injection of Rh immunoglobulin during the delivery episode.
	Date postpartum Rh immunoglobulin – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day of the postpartum injection of Rh immunoglobulin during the delivery episode.
	Drugs received during delivery admission – Antihypertensives	A flag (i.e., Yes, Null) to indicate mother received antihypertensive medication during the delivery episode of care.
	Drugs received during delivery admission – Steroids for lung maturation	A flag (i.e., Yes, Null) to indicate mother received steroid medication during the inpatient delivery episode of care, or for transport.

08/09 onwards
)8/09 onwards
08/09 onwards
08/09-onwards
800 800 800 800 800 800 800 800 800

	Postpartum urinary tract infection agent – Infectious agent 2	Other infectious agent causing positive maternal urine culture results during the postpartum period.	2008/09-onwards
	Postpartum positive blood culture	A flag (i.e., Yes, Null) to indicate mother's blood culture test results are positive.	2008/09-onwards
	Postpartum positive blood culture agent – Infectious Agent 1	Most significant infectious agent causing positive maternal blood culture results during the postpartum period.	2008/09-onwards
	Postpartum positive blood culture agent – Infectious Agent 2	Other infectious agent causing positive maternal blood culture results during the postpartum period.	2008/09-onwards
	Postpartum positive other culture	A flag (i.e., Yes, Null) to indicate lab culture results were positive (other than urine or blood).	2008/09-onwards
	Postpartum positive other culture agent – Infectious agent 1	Most significant infectious agent from other maternal positive culture results during the postpartum period.	2008/09-onwards
	Postpartum positive other culture agent - Infectious agent 2	Other infectious agent from other maternal positive culture results during the postpartum period.	2008/09-onwards
	Main Patient Service	Categorizes mothers according to related diseases, conditions and treatments.	
Diag	nosis and procedures		
	Diagnosis Prefix	Alphanumeric character to further define a diagnosis code.	
	Diagnosis Code	Medical diagnostic code reflecting the diagnosis or condition of mother while in hospital. Note: International Classification of Diseases - ICD 9 - Assigned from April 1, 2000 to March 31, 2004. ICD-10-CA - Assigned starting with April 1, 2004 discharges using ICD-10-CA v2003, v2006, v2009, or v2012 (as applicable).	
	Diagnosis Type	Diagnosis type corresponding with the diagnosis codes (e.g., most responsible diagnosis, pre-admit comorbidity, secondary diagnosis, etc.)	
	Procedure Code	Code(s) for procedures performed during the episode of care. Note: Canadian Classification of Diagnostic, Therapeutic and Surgical Procedures (CCP) - Assigned from April 1, 2000 to March 31, 2004 discharges. Canadian Classification of Health Interventions (CCI) - Assigned starting with April 1, 2004 discharges using CCI v2003, v2006, v2009, or v2012 (as applicable).	
	Procedure status	Procedure status attribute.	2004/05-onwards
	Procedure location	Procedure anatomical location.	2004/05-onwards
	Procedure extent	Procedure extent.	2004/05-onwards
	Procedure Date – Year	Year of the procedure.	
	Procedure Date – Month	Month of the procedure.	
	Procedure Date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day of the procedure.	
	Procedure doctor service	The procedure provider service.	
	Anesthetic agent for procedure	Type of anesthesia used for the procedure (e.g., local, epidural, spinal, etc).	

Mother Postpartum Transfer/Readmission Episode of Care Information (patient discharges from April 1, 2008 to March 31, 2018) Place of Mother's usual residence - HA - **Research rationale describing why this field is required must be supplied in your application before it will be considered for release: The Health Authority (HA) of mother's usual residence as determined by resident postal code. Click here to enter text. Mother's date of birth - Year Mother's date of birth - Month Mother's date of birth – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers Place of postpartum admission - Replaced by project-Location where mother received care. specific identification number Institution From - Replaced by project-specific Institution from which mother arrived to the current identification number episode of care. A flag (i.e., Yes, Null) to indicate mother was Mother transferred in transferred in from another acute care institution for the current episode of care. Institution To - Replaced by project-specific Institution to which mother was transferred from the identification number current episode of care. A flag (i.e., Yes, Null) to indicate mother was transferred out to another institution from the current Mother transferred out episode of care. Year mother was admitted for the current episode of Admission Date - Year Month mother was admitted for the current episode Admission Date - Month of care. Admission Date - Day - This field is available for use in Day mother was admitted for the current episode of analysis by the WHRI Data Analyst, but will not be care. provided to researchers Year mother was discharged from the current Discharge Date - Year episode of care. Month mother was discharged from the current Discharge Date - Month episode of care. Discharge Date - Day - This field is available for use in Day mother was discharged from the current episode analysis by the WHRI Data Analyst, but will not be of care. provided to researchers The fiscal year during which the mother was Fiscal year discharged (i.e., Apr 1 to Mar 31). Total Length of Stay (hours) Length of stay of admission expressed in hours. Delivery date - Year Year the woman delivered. Delivery date - Month Month the woman delivered. Delivery date - Day - This field is available for use in analysis by the WHRI Data Analyst, but will not be Day the woman delivered. provided to researchers

	Place of delivery – Replaced by project-specific identification number	Institution where the mother delivered.			
Othe	Other episode of care information				
	Blood transfusion given	A flag (i.e., Yes, Null) to indicate mother received whole or packed red blood cells during this admission.			
	Blood transfusion units – Number of units transfused postpartum	Total number of units of whole or packed red blood cells the mother received during the episode of care.			
	Health care provider(s) service	Provider's specialty service number.			
	Health care provider(s) type	Health care provider's role in the care of the mother during episode of care (e.g., most responsible, resident/intern, allied health etc.)			
Post	delivery information				
	HELLP Syndrome	A flag (i.e., Yes, Null) to indicate mother was diagnosed with HELLP Syndrome			
	Acute Fatty Liver	A flag (i.e., Yes, Null) to indicate mother diagnosed with acute fatty liver during current pregnancy or postpartum period.			
	Liver hematoma	A flag (i.e., Yes, Null) to indicate mother diagnosed with liver hematoma during current pregnancy or postpartum period.			
	Postpartum Special Care Unit Days	Number of days mother spent in any Special Care Unit (ICU, CCU, etc.).			
	Postpartum hemoglobin date – Year	Year of mother's lowest postpartum hemoglobin result during episode of care.			
	Postpartum hemoglobin date – Month	Month of mother's lowest postpartum hemoglobin result during episode of care.			
	Postpartum hemoglobin date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day of mother's lowest postpartum hemoglobin result during episode of care.			
	Postpartum hemoglobin value	Value of postpartum hemoglobin test result during episode of care.			
	Postpartum infection	Mother had an infection during the episode of care (i.e., Yes, No, Unknown).			
	Postpartum wound infection	A flag (i.e., Yes, Null) to indicate mother had a postpartum wound infection.			
	Postpartum wound infection – Type	Specific location of the mother's postpartum wound infection.			
	Postpartum wound infection – Severity	Degree of the mother's postpartum wound infection.			
	Postpartum urinary tract infection – Type	Type of postpartum urinary tract infection			
	Postpartum urinary tract infection agent – Infectious agent 1	Most significant infectious agent causing positive maternal urine culture results during the episode of care.			
	Postpartum urinary tract infection agent – Infectious agent 2	Other infectious agent causing positive maternal urine culture results during the episode of care.			
	Postpartum positive blood culture	A flag (i.e., Yes, Null) to indicate mother's blood culture test results are positive.			
	Postpartum positive blood culture agent – Infectious Agent 1	Most significant infectious agent causing positive maternal blood culture results during the episode of care.			
	Postpartum positive blood culture agent – Infectious Agent 2	Other infectious agent causing positive maternal blood culture results during the episode of care.			

	Postpartum positive other culture	A flag (i.e., Yes, Null) to indicate lab culture results were positive (other than urine or blood).	
	Postpartum positive other culture agent – Infectious agent 1	Most significant infectious agent from other maternal positive culture results during the episode of care.	
	Postpartum positive other culture agent - Infectious agent 2	Other infectious agent from other maternal positive culture results during the episode of care.	
	Main Patient Service	Categorizes mothers according to related diseases, conditions and treatments.	
Diag	noses and procedures		
	Diagnosis Prefix	Alphanumeric character to further define a diagnosis code.	
	Diagnosis Code	Medical diagnostic code reflecting the diagnosis or condition of mother while in hospital. Note: ICD-10-CA - Assigned starting with April 1, 2008 discharges using ICD-10-CA v2006, v2009, or v2012 (as applicable).	
	Diagnosis Type	Diagnosis type corresponding with the diagnosis codes (e.g., most responsible diagnosis, pre-admit comorbidity, secondary diagnosis, etc)	
	Procedure Code	Code(s) for procedures performed during the episode of care. Note: Canadian Classification of Health Interventions (CCI) - Assigned starting April 1, 2008 discharges using CCI v2006, v2009, or v2012 (as applicable).	
	Procedure status	Procedure status attribute	
	Procedure location	Procedure anatomical location	
	Procedure extent	Procedure extent	
	Procedure Date – Year	Year of the procedure.	
	Procedure Date – Month	Month of the procedure.	
	Procedure Date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day of the procedure.	
	Procedure doctor service	The procedure provider service.	
	Anesthetic agent for procedure	Type of anesthesia used for the procedure (e.g., general, spinal, epidural, etc.).	
Baby Newborn Episode of Care Information (April 1, 2000 to March 31, 2018)			
Baby Newborn records include births in acute care facilities and births at home attended by registered midwives. NOTE: Until March 31, 2014, a baby born at home under the care of a registered midwife who was admitted to acute care within 24 hours of birth will have the acute care admission as the Baby Newborn record (total record count=1). Effective April 1, 2014, a baby born at home with a registered midwife and who was admitted to acute care within 24 hours of birth will have a Baby Newborn record at home and a subsequent Baby Transfer/Readmission acute care record (total record count=2).			
	Place of baby's usual residence – HA – **Research rationale describing why this field is required must be supplied in your application before it will be considered for release: Click here to enter text.	The Health Authority (HA) of baby's usual residence as determined by resident postal code	
		l	

	Baby Sequence	The incremental sequence number of babies born from the <i>current</i> pregnancy (e.g. twin A = sequence 1, twin B = sequence 2).	
	Number of births	The total number of babies delivered from the current pregnancy.	
	Baby Date of Birth – Year	Year baby was born.	
	Baby Date of Birth – Month	Month baby was born.	
	Baby Date of Birth – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day baby was born.	
	Sex	Biological sex of the newborn.	
Curr	ent admission information		
	Place of birth – Replaced by project-specific identification number	Location where baby received care.	
	Institution To – Replaced by project-specific identification number	Institution to which baby was transferred from the current episode of care.	
	Transfer up (to higher level of care)	A flag (i.e., Yes, Null) to indicate newborn transferred to a hospital with a higher level of care directly from the birth episode.	
	Admission Date – Year	Year baby was admitted to the current episode of care.	
	Admission Date – Month	Month baby was admitted to the current episode of care.	
	Admission Date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day baby was admitted to the current episode of care.	
	Discharge Date – Year	Year baby was discharged from the current episode of care.	
	Discharge Date – Month	Month baby was discharged from the current episode of care.	
	Discharge Date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day baby was discharged from the current episode of care.	
	Fiscal Year	The fiscal year during which the baby was discharged (i.e., Apr 1 to Mar 31).	
	Length of Stay (Hours)	Baby's length of stay for admission expressed in hours. Note: Always null for home births.	
	Neonatal Intensive Care Unit days (Level II)	Total number of days baby was in Neonatal Intensive Care Unit Level II. Note: Changes over time to calculation method. Also, documented data quality issues from 2010/11 onwards.	2004/05 onwards
	Neonatal Intensive Care Unit days (Level III)	Total number of days baby was in Neonatal Intensive Care Unit Level III. Note: Changes over time to calculation method. Also, documented data quality issues from 2010/11 onwards.	2004/05 onwards
	Admission weight	Admission weight in grams.	
	Discharge weight	Baby's weight (in grams) at discharge.	
\boxtimes	Gestational age at birth by newborn exam	Baby's gestational age (in completed weeks) based on care provider's physical assessment and neuromuscular assessment of the newborn at birth. Note: Checked automatically because this field	

		should always be requested when delivery or		
		newborn records requested.		
	Gestational age at birth from maternal chart	Baby's gestational age (in completed weeks) documented by the care provider before delivery, determined by maternal last menstrual period and/or ultrasound. Note: Checked automatically because this field should always be requested when delivery or newborn records requested.		
	Gestational age at birth, in completed weeks – calculated by algorithm incorporating LNMP, first U/S, newborn examination, and maternal chart	Gestational age at birth, in completed weeks – calculated by algorithm incorporating LNMP, first U/S, newborn examination, and maternal chart. Note: Checked automatically because this field should always be requested when delivery or newborn records requested.		
	Birth length	Length of baby at birth (in centimeters).		
	Birth head circumference	Head circumference of baby at birth (in centimeters).		
	Birth Type	Identifies birth type for births at or after 20 weeks gestation or weighing at least 500 grams (i.e., stillbirth, live birth).		
	Stillbirth timing	The stage in labour when the stillbirth occurred (e.g., stillbirth after onset of labour, stillbirth prior to onset of labour).		
	Vitamin K	Newborn received vitamin K (i.e., Yes, No, Unknown).	2000/01-2007/08	
	Eye prophylaxis given	Baby received erythromycin or other eye prophylaxis (i.e., Yes, No, Unknown).	2000/01-2007/08	
	Breastfeeding at discharge	Indicates if mother is breastfeeding the baby at discharge (i.e., Yes, No, Unknown).	2000/01-2003/04	
	Newborn feeding	The type of feeding given to the newborn during the entire hospital stay, including discharge (e.g., Exclusive breast milk, breast milk and formula, formula, etc.).	2004/05-onwards	
	Breast feeding initiation	Time frame during which breastfeeding first commenced/attempted following delivery, regardless of whether the baby latched.	2008/09-onwards	
	Health care provider(s) service	Provider's specialty service number.		
	Health care provider(s) type	Health care provider's role in the care of the baby during episode of care (e.g., most responsible, resident/intern, allied health etc.)		
	Discharged to	Where the baby was discharged to, or the status of the baby at the time of discharge (e.g., adoption, death/stillbirth, foster home, home, other hospital, unknown). Records with Adoption and Foster home are		
	Additional and the same field to	excluded from the dataset Value of first temperature taken within the first hour	2000/00	
	1st temperature within 1st hour after birth	of birth (in Celsius to 1 decimal place). A flag (i.e., Yes, Null) to indicate surfactant	2008/09-onwards	
	Surfactant Given	administered during hospital admission.	2008/09-onwards	
	Antibiotics Given	A flag (i.e., Yes, Null) to indicate antibiotics were administered during hospital admission.	2008/09-onwards	
Birth	Birth information			
	Apgar 1 minute	Apgar score at 1 minute.		

Apgar 5 minutes	Apgar score at 5 minutes.	
Apgar 10 minutes	Apgar score at 10 minutes.	
Meconium thick	A flag (i.e., Yes, Null) to indicate meconium described as thick or particulate at birth.	2000/01-2003/04
Meconium	A flag (i.e., Yes, Null) to indicate presence of thick or thin meconium at any time during the intrapartum period, including delivery.	2004/05-onwards
Drugs for resuscitation / stabilization	Administration of medication to the newborn for resuscitative/stabilization purposes during the birth episode (i.e., Yes, No, Unknown).	
Suction – Perineum	A flag (i.e., Yes, Null) to indicate baby is suctioned at the perineum upon delivery of the head.	2000/01-2007/08
Suction – Oropharynx	A flag (i.e., Yes, Null) to indicate clearing of the newborn's airway at the level of the oropharynx.	
Suction – Trachea	A flag (i.e., Yes, Null) to indicate clearing of the newborn's airway at the level of the trachea.	
Suction – Unspecified	A flag (i.e., Yes, Null) to indicate clearing of the newborn's airway at an unspecified level.	
Oxygen for resuscitation	A flag (i.e., Yes, Null) to indicate baby received oxygen for immediate resuscitation.	
Oxygen for resuscitation – Age started	Age in minutes when oxygen for resuscitation started.	
Oxygen for resuscitation – Age stopped	Age in minutes when oxygen for resuscitation ended.	
Total length of time oxygen given for resuscitation	Total minutes baby received oxygen for immediate resuscitation.	
IPPV mask given for resuscitation	A flag (i.e., Yes, Null) to indicate newborn received intermittent positive pressure ventilation (IPPV) for immediate resuscitation via mask.	
IPPV mask given for resuscitation – Age started	Age in minutes when intermittent positive pressure ventilation (IPPV) by mask started.	
IPPV mask given for resuscitation – Age stopped	Age in minutes when intermittent positive pressure ventilation (IPPV) by mask for resuscitation ended.	
Total length of time IPPV mask given for resuscitation	Total minutes baby received intermittent positive pressure ventilation (IPPV) by mask for immediate resuscitation.	
IPPV ETT given for resuscitation	A flag (i.e., Yes, Null) to indicate newborn received intermittent positive pressure ventilation (IPPV) for immediate resuscitation via endotracheal tube (ETT).	
IPPV ETT given for resuscitation – Age started	Age in minutes when intermittent positive pressure ventilation (IPPV) by endotracheal tube (ETT) started.	
IPPV ETT given for resuscitation – Age stopped	Age in minutes when intermittent positive pressure ventilation (IPPV) by endotracheal tube (ETT) for resuscitation ended	
Total length of time IPPV ETT given for resuscitation	Total minutes baby received intermittent positive pressure ventilation (IPPV) by endotracheal tube (ETT) for immediate resuscitation.	
Chest compressions given for resuscitation	A flag (i.e., Yes, Null) to indicate baby received chest compressions for immediate resuscitation.	
Chest compressions given for resuscitation – Age Started	Age in minutes when chest compressions for resuscitation started	
Chest compressions given for resuscitation – Age Stopped	Age in minutes when chest compressions for resuscitation ended	
Total length of time compressions given for resuscitation	Total minutes baby received chest compressions for immediate resuscitation.	

	Total ventilator days	Total number of days (in whole numbers) baby was on a ventilator.	
	Total CPAP days	Total number of days (in whole numbers) baby was on Continuous Positive Airway Pressure (CPAP).	2008/09-onwards
	Total oxygen days	Total number of days (in whole numbers) baby received continuous oxygen therapy or nasal prongs.	
	Total TPN days	Total number of days (in whole numbers) the baby received any total parenteral nutrition (TPN).	
	Cord arterial gases pH	pH value of cord arterial blood gases, obtained from the umbilical artery.	
	Cord arterial gases base excess/deficit	Base excess (+) or deficit (-) value of the cord arterial blood gases, obtained from the umbilical artery.	
	Positive Blood Culture	A flag (i.e., Yes, Null) to indicate baby's blood culture test results were positive.	2008/09-onwards
	Positive blood culture agent – Infectious Agent 1	Most significant infectious agent causing positive blood culture results in the baby.	2008/09-onwards
	Positive blood culture agent – Infectious Agent 2	Other infectious agent causing positive blood culture results in the baby.	2008/09-onwards
	Positive urine culture	A flag (i.e., Yes, Null) to indicate baby's urine culture test results were positive.	2008/09-onwards
	Positive urine culture – Infectious Agent 1	Most significant infectious agent causing positive urine culture results in the baby.	2008/09-onwards
	Positive urine culture – Infectious Agent 2	Other infectious agent causing positive urine culture results in the baby.	2008/09-onwards
	Positive other culture	A flag (i.e., Yes, Null) to indicate lab culture test results were positive for the baby (other than blood and urine).	2008/09-onwards
	Positive other culture – Infectious Agent 1	Most significant infectious agent causing other positive culture results in the baby, other than blood and urine.	2008/09-onwards
	Positive other culture – Infectious Agent 2	Other infectious agent causing other positive culture results in the baby, other than blood and urine.	2008/09-onwards
	Main Patient Service	Categorizes babies according to related diseases, conditions and treatments.	
Diaç	gnosis and procedures		
	Diagnosis prefix	Alphanumeric character to further define a diagnosis code.	
	Diagnosis Code	Medical diagnostic code reflecting the diagnosis or condition of mother while in hospital. Note: International Classification of Diseases - ICD 9 - Assigned from April 1, 2000 to March 31, 2004. ICD-10-CA - Assigned starting with April 1, 2004 using ICD-10-CA v2003, v2006, v2009, or v2012 (as applicable).	
	Diagnosis Type	Diagnosis type corresponding with the diagnosis codes (e.g., most responsible diagnosis, pre-admit comorbidity, secondary diagnosis, etc)	
	Procedure Code	Code(s) for procedures performed during the episode of care. Note: Canadian Classification of Diagnostic, Therapeutic and Surgical Procedures - Assigned from April 1, 2000 to March 31, 2004. Canadian Classification of Health Interventions (CCI) - Assigned effective April 1, 2004 using CCI v2003, v2006, v2009, or v2012 (as applicable).	
	Procedure status	Procedure status attribute	2004/05-onwards
	Procedure location	Procedure anatomical location	2004/05-onwards

Procedure extent	Procedure extent	2004/05-onwards
Procedure Date – Year	Year of the procedure.	
Procedure Date – Month	Month of the procedure.	
Procedure Date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day of the procedure.	
Procedure doctor service	The procedure provider service.	
Anesthetic agent for procedure	Type of anesthesia used for the procedure (e.g., local, epidural, spinal etc.).	
y Transfer/Readmission Episode of Care I ril 1, 2000 to March 31, 2018)	nformation	
Place of baby's usual residence – HA – **Research rationale describing why this field is required must be supplied in your application before it will be considered for release: Click here to enter text.	The Health Authority (HA) of baby's usual residence as determined by resident postal code	
Baby Sequence	The incremental sequence number of babies born from the <i>current</i> pregnancy (e.g. twin A = sequence 1, twin B = sequence 2).	
Number of births	The total number of babies delivered from the current pregnancy.	
Sex	Biological sex of the newborn.	
Place of admission – Replaced by project-specific identification number	Location where baby received care	
Institution from – Replaced by project-specific identification number	Institution from which baby was admitted to the current episode of care.	
Institution To – Replaced by project-specific identification number	Institution to which baby was transferred from the current episode of care.	
Admission Date – Year	Year baby was admitted to the current episode of care.	
Admission Date – Month	Month baby was admitted to the current episode of care.	
Admission Date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day baby was admitted to the current episode of care.	
Discharge Date – Year	Year baby was discharged from the current episode of care.	
Discharge Date – Month	Month baby was discharged from the current episode of care.	
Discharge Date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day baby was discharged from the current episode of care.	

Fiscal Year	The fiscal year during which the baby was discharged (i.e., Apr 1 to Mar 31).	
Length of Stay (Hours)	Baby's length of stay for admission expressed in hours.	
Neonatal Intensive Care Unit days (Level II)	Total number of days baby was in Neonatal Intensive Care Unit Level II. Note: Changes over time to calculation method. Also, documented data quality issues from 2010/11 onwards.	2004/05-onwards
Neonatal Intensive Care Unit days (Level III)	Total number of days baby was in Neonatal Intensive Care Unit Level III. Note: Changes over time to calculation method. Also, documented data quality issues from 2010/11 onwards.	2004/05-onwards
Admission weight	Admission weight in grams.	
Discharge weight	Baby's weight (in grams) at discharge.	
Discharged to	Where the baby was discharged to, or the status of the baby at the time of discharge (i.e., adoption, death/stillbirth, foster home, home, other hospital, unknown). Records with Adoption and Foster home are	
Health care provider(s) service	excluded from the dataset Provider's specialty service number.	
 Treatur care provider(s) service	Health care provider's role in the care of the baby	
Health care provider(s) type	during episode of care (e.g., most responsible, resident/intern, allied health).	
1st temperature within 1st hour after birth	Value of first temperature taken within the first hour of birth (in Celsius to 1 decimal place).	2008/09-onwards
Surfactant Given	A flag (i.e., Yes, Null) to indicate surfactant administered during hospital admission.	2008/09-onwards
Antibiotics Given	A flag (i.e., Yes, Null) to indicate antibiotics were administered during hospital admission.	2008/09-onwards
Total ventilator days	Total number of days (in whole numbers) baby was on a ventilator.	
Total CPAP days	Total number of days (in whole numbers) baby was on Continuous Positive Airway Pressure (CPAP).	2008/09-onwards
Total oxygen days	Total number of days (in whole numbers) baby received continuous oxygen therapy or nasal prongs.	
Total TPN days	Total number of days (in whole numbers) the baby received any total parenteral nutrition (TPN).	
Positive Blood Culture	A flag (i.e., Yes, Null) to indicate baby's blood culture test results were positive.	2008/09-onwards
Positive blood culture agent – Infectious Agent 1	Most significant infectious agent causing positive blood culture results in the baby.	2008/09-onwards
Positive blood culture agent – Infectious Agent 2	Other infectious agent causing positive blood culture results in the baby.	2008/09-onwards
Positive urine culture	A flag (i.e., Yes, Null) to indicate baby's urine culture test results were positive.	2008/09-onwards
Positive urine culture – Infectious Agent 1	Most significant infectious agent causing positive urine culture results in the baby.	2008/09-onwards
Positive urine culture – Infectious Agent 2	Other infectious agent causing positive urine culture results in the baby.	2008/09-onwards
Positive other culture	Lab culture test results were positive for the baby (other than blood and urine).	2008/09-onwards
Positive other culture – Infectious Agent 1	Most significant infectious agent causing other positive culture results in the baby, other than blood and urine.	2008/09-onwards

Positive other culture – Infectious Agent 2	Other infectious agent causing other positive culture results in the baby, other than blood and urine.	2008/09-onwards
Main Patient Service	Categorizes babies according to related diseases, conditions and treatments.	
Diagnoses and procedures		
Diagnosis prefix	Alphanumeric character to further define a diagnosis code.	
Diagnosis Code	Medical diagnostic code reflecting the diagnosis or condition of mother while in hospital. Note: International Classification of Diseases - ICD 9 - Assigned from April 1, 2000 to March 31, 2004. ICD-10-CA - Assigned effective April 1, 2004 using ICD-10-CA v2003, v2006, v2009, or v2012 (as applicable).	
Diagnosis Type	Diagnosis type corresponding with the diagnosis codes (e.g., most responsible diagnosis, pre-admit comorbidity, secondary diagnosis, etc)	
Procedure Code	Code(s) for procedures performed during the episode of care. Note: Canadian Classification of Diagnostic, Therapeutic and Surgical Procedures - Assigned from April 1, 2000 to March 31, 2004. Canadian Classification of Health Interventions (CCI) - Assigned effective April 1, 2004 using CCI v2003, v2006, v2009, or v2012 (as applicable).	
Procedure status	Procedure status attribute	2004/05 onwards
Procedure location	Procedure anatomical location	2004/05 onwards
Procedure extent	Procedure extent	2004/05 onwards
Procedure Date – Year	Year of the procedure.	
Procedure Date – Month	Month of the procedure.	
Procedure Date – Day – This field is available for use in analysis by the WHRI Data Analyst, but will not be provided to researchers	Day of the procedure.	
Procedure doctor service	The procedure provider service.	
Anesthetic agent for procedure	Type of anesthesia used for the procedure (e.g., local, epidural, spinal etc.).	